



ISSN: 3062-3405

Annals of Orthodontics and Periodontics Specialty

Volume 4, Page No: 171-180

Available Online at: aopsj.com

Original Article

Clinical Evaluation of Laser-Assisted New Attachment Procedure® (LANAP®) Surgical Treatment of Chronic Periodontitis: A Retrospective Case Series of 1-Year Results in 22 Consecutive Patients

Alexandra Micu^{1*}, Corina Nour¹, Gheorghe Szilagyi¹

1. Faculty of Dental Medicine, George Emil Palade University of Medicine, Pharmacy, Science, and Technology of Targu Mures, 38 Gh. Marinescu Str., 540142 Targu Mures, Romania.

*E-mail ✉ Alexandermicu@outlook.com

Abstract

Treatment for periodontitis has evolved over the years as new technologies have become available. Currently, lasers seem attractive as a treatment modality, but their effectiveness needs to be verified. The purpose of this project was to evaluate Laser Assisted New Attachment Procedure® (LANAP®) surgery as a single treatment modality. As part of a mandatory training program for periodontists and other dentists, 22 consecutive patients diagnosed with moderate to severe periodontitis (probing depth [PD] up to 11 mm) were treated with the LANAP® surgical approach using a 1064-nm Nd:YAG laser as part of a multi-step protocol. Following single-session active therapy, they were entered into a maintenance program. Their clinical status was re-evaluated at 12–18 months following surgery. All 22 patients completed the 12- to 18-month follow-up. PD, clinical attachment level, and furcation (FURC) showed substantial improvement. Recession was minimal (mean, 0.1 mm), while 93.5% of PD measurements were 3 mm or less at re-evaluation. Furthermore, 40% of grade 2 FURC closed clinically. Within the limits of this case series, LANAP® was found to be an effective, minimally invasive, laser surgical therapy for moderate to advanced periodontitis.

Key words: Laser, Periodontitis, Surgery, LANAP®

How to cite this article: Micu A, Nour C, Szilagyi G. Clinical Evaluation of Laser-Assisted New Attachment Procedure® (LANAP®) Surgical Treatment of Chronic Periodontitis: A Retrospective Case Series of 1-Year Results in 22 Consecutive Patients. Ann Orthod Periodontics Spec. 2024;4:171-80. <https://doi.org/10.51847/zFaFw46b>

Received: 27 February 2024; Revised: 07 June 2024; Accepted: 09 June 2024

Introduction

Periodontitis is a major dental disease, especially in adults. It has been reported that overall about 42% of U.S. adults have periodontitis, with about 34% having moderate to severe cases [1]. Historically, a variety of non-surgical and surgical treatments have been used to reduce probing depth (PD) and inflammation. While non-surgical treatments are modestly effective [2-6], surgical treatment is often recommended as a more effective modality for the problems of increased PD, loss of clinical attachment level (CAL), and bone loss; in recent years, studies have focused on the regeneration of lost or damaged supporting tissues [7, 8]. Since most patients would like to keep their teeth, the main goal of all dental therapies is the retention of natural teeth in health, comfort, function, and esthetics. Clinical parameters such as PD, CAL, non-bleeding percentage (nBOP), and the modified gingival index (mGI), are surrogate markers for periodontal status, but provide practical clinical



© 2024 Annals of Orthodontics and Periodontics Specialty

Open Access – This Article is licensed under CC BY NC SA 4.0. To view a copy of this license, visit <https://creativecommons.org/licenses/by-nc-sa/4.0/>

information. Generally, the treatment goals are to have no or minimal gingival inflammation or bleeding, shallow (3 mm or less) PDs, and minimal gingival recession (REC).

Lasers have been used as a single treatment, before or after scaling and root planing, and combined with surgical treatment. While non-surgical laser treatments have demonstrated variable results [9], a definitive laser surgical approach [8, 10], Laser-Assisted New Attachment Procedure® (LANAP®), has shown promise in published case reports as a single treatment, laser based, surgical approach to improving periodontal parameters [11-16].

Information regarding the results of different treatment modalities is continually needed in periodontics. This is particularly true of newer treatment approaches and techniques, such as LANAP®, which uses a particular laser wavelength (Nd:YAG, 1,064 nm). LANAP® can only be legally performed after required training, including hands-on work with actual patients, is completed. Details about the protocol can be found in articles by Aoki *et al.* [10] and Jha *et al.* [17]. Human histologic evidence supports the concept of periodontal regeneration/new attachment with this procedure [18, 19]. The LANAP® protocol includes active, definitive full-mouth surgical treatment as the initial therapy, and regular follow-up periodontal maintenance and occlusal adjustments, including a biteguard. This protocol saves patient treatment time, generally reduces patient costs, and potentially has equal or better results than the typical 2 step approach (initial therapy followed by surgical therapy) in achieving the treatment goals stated above. The typical treatment and follow-up scheme for LANAP® patients is shown in **Table 1**.

Table 1. LANAP® treatment protocol

LANAP® treatment protocol		
Patient name:	Date:	
Time interval date:	Time needed	Procedure
1st appt.	1 hour	Periodontal charting (DDS)
		X-rays (DA)
		Treatment plan consultation (DDS)
10–14 days	30 minutes	Prescriptions
		Preoperative and postoperative Information
		LANAP® surgery, half mouth
ASAP	2 1/2 hours	Postoperative (DDS)
		Occlusal adjustment first side
		LANAP® surgery, other half mouth
7–12 days	2 1/2 hours	Postoperative - (DDS)
		Occlusal adjustment - (DDS)
		RDH polish
10–14 days	30 minutes	Possible occlusal adjustment - (DDS)
		RDH polish
		Possible occlusal adjustment - (DDS)
14–28 days	30 minutes	Impressions for biteguard - (DA)
		RDH polish
		Possible occlusal adjustment - (DDS)
1 month	30 minutes	RDH polish
		Possible occlusal adjustment - (DDS)
		Biteguard delivery - (DDS)
1 month	30 minutes	Periodontal maintenance (RDH)
		Possible occlusal adjustment - (DDS)
		Periodontal maintenance (RDH)
3 months	1 hour	Possible occlusal adjustment - (DDS)
		Periodontal maintenance (RDH)
		Possible occlusal adjustment - (DDS)

3 months	1 hour	Periodontal maintenance (RDH)
		Possible occlusal adjustment - (DDS)
3 months	1 hour	Periodontal maintenance (RDH)
		Possible occlusal adjustment - (DDS)
2–3 weeks	1 hour	Periodontal charting - (DDS)
		X-rays - (DA)
Q3 months	1 hour	Periodontal maintenance (RDH)
		Possible occlusal adjustment - (DDS)
Annually	1 hour	Periodontal charting - (DDS)
		X-rays (DA)

LANAP®: Laser-Assisted New Attachment Procedure®, DDS: doctor of dental surgery; DA: dental assistant, RDH: registered dental hygienist.

The purpose of this retrospective case series is to report clinical results following LANAP® treatment. The radiographic results will be reported separately.

Materials and Methods

The current evaluation documents the clinical periodontal changes following LANAP® in 22 consecutively treated patients after 12–18 months of follow-up. This retrospective study was accorded exempt status after review by the Western Institutional Review Board due to its retrospective nature, so there was no need for separate informed consent (WIRB #281385). The physical and digital records of the patients who met the predetermined eligibility criteria were de-identified using a computer-generated numbering scheme.

Patients presenting to the University of Colorado School of Dental Medicine Graduate Periodontics Clinic diagnosed by an experienced clinician with generalized moderate to severe chronic periodontitis (i.e., meeting the case definition of generalized moderate to severe chronic periodontitis used at that time (American Dental Association [ADA] case type 4) (Armitage [20]), were treated with LANAP® if they met the following inclusion criteria: generalized moderate to severe chronic periodontitis, no definitive periodontal treatment nor antibiotic use in the last 12 months, no systemic diseases that would influence the results of periodontal therapy, and no pregnancy or breastfeeding. Smokers were included. The study period was from November 2006 to November 2010.

Patients had complete periodontal charting and 7 vertical bitewing and panoramic radiographs completed at baseline and at 12–18 months after active treatment. Clinical and radiographic findings, the diagnosis, etiologies, prognosis and all primary treatment options were discussed with the patients at a separate treatment planning appointment. These included no treatment, scaling and root planing, flap surgery (either subtractive or additive), laser surgery, and/or extraction and replacement. All patients expressed a desire to keep their teeth if possible, and opted and consented orally and in writing for laser surgery. They were advised that their treatment would be performed by “trainees” using the laser under supervision.

No initial periodontal treatment was provided. Patients who consented to LANAP® went directly to laser surgery. LANAP® was performed in 1 half mouth at a time using local anesthesia. The patients were treated with LANAP® during a 3-day training program developed by the Institute for Advanced Laser Dentistry (IALD). Patient treatment was supervised one-on-one by various certified and calibrated instructors of the IALD, with overall on-site supervision by the author.

Clinical periodontal measurements were performed by the author, an American Board of Periodontology–certified periodontist, at baseline and at 12–18 months after completion of the second half mouth of full-mouth LANAP®. Standard parameters were evaluated including PD; the free gingival margin (FGM) level from the cemento-enamel junction; CAL, was calculated from the prior 2 measurements; bleeding on probing, reported as the nBOP (bleeding on probing was deemed positive if bleeding occurred within 30 seconds after probing, and the score was converted to a non-bleeding value); furcation (FURC) involvement [21]; tooth mobility (MOB) [22]; the modified plaque index (mPI) [23]; and the mGI [24]. The investigator was calibrated with measurements in 3 patients who were diagnosed as having chronic periodontitis and were

not included in this study. FGM position, PD, nBOP, and FURC were repeated after a 2-day interval. The intra-examiner kappa scores were 0.94 for PD, nBOP, and FURC, and 0.88 for FGM and mPI. FGM, PD, nBOP, and mPI were recorded at 6 sites of all teeth present except third molars using a manual periodontal probe (UNC15, Hu-Friedy, Inc., Chicago, IL, USA). FURC measurements were made with a calibrated Nabers probe (Hu-Friedy, Inc.).

During the laser application, laser safety rules were followed, including wavelength-specific eyewear protection for all individuals in the operatory, close monitoring of laser settings, and careful observation of the tissue response to the laser energy.

Using a 1064-nm PerioLase MPV-7™ laser, Millennium Dental Technologies, Cerritos, CA, USA **Figure 1**, the settings were as follows: 3.6 W average power, 20 Hz pulse repetition rate, pulse duration of 100 μ s for first laser application and 550 μ s for the second laser application delivered through a 360-micron quartz fiber in a special handpiece. This yielded an energy delivery of 180 mJ per pulse, an energy density of 177 J/cm², and a power density of 3,537 W/cm². The peak power was 1800 W during the first pass and 327 W during the second laser pass. The goal was to deliver 10–12 J/ per millimeter of PD, based on a recommended maximum of 12–16 J per millimeter of PD, which was almost always achieved (data on file) [25].



Figure 1. PerioLase MVP-7 Nd:YAG laser, Millennium Dental Technologies, Inc.

Between the 2 laser applications, the roots were thoroughly debrided with a piezoelectric ultrasonic scaler and tips (Piezon 400, Electro Medical Systems, Nyon, CHE); and decortication/intra-marrow penetration was performed at the base and along the walls of the bony defects [26]. In all cases, in accordance with LANAP®, mobility and other manifestations of occlusal pathology were assessed. The occlusion was carefully addressed and managed. No additive materials (such as bone replacement grafts, membranes, or biologics), sutures, or periodontal dressings were used.

Post-surgically, patients were prescribed ibuprofen (800 mg) for possible pain, doxycycline (100 mg, twice daily for 7 days) for residual bacteria, and a 0.12% chlorhexidine rinse to control new bacterial accumulation, since mechanical plaque removal was to be avoided on each treated side for the first 2–3 weeks.

Following surgical treatment of the second side, patients were seen frequently for general evaluations, oral hygiene instructions, supra-gingival polishing/de-plaquing to minimize inflammation, and occlusal adjustment to minimize unfavorable forces. After 4–6 weeks, a maxillary flat plane biteguard was delivered. Periodontal maintenance every 3 months followed. Occlusal evaluation and adjustment were planned for each visit (**Table 1**). As per protocol, neither probing nor sub-gingival instrumentation was performed unless a clinical need such as evident calculus or inflammation arose (none did) until the 12- to 18-month re-evaluation.

Results and Discussion

All 22 patients completed active treatment and the 12- to 18-month follow-up sessions, except for isolated instances (no more than 2) of missed appointments for any given patient.

Table 2 presents the age, sex, ethnicity, tooth status and smoking habit of the patients. The initial sextant diagnosis was generalized moderate to severe chronic periodontitis in 18 cases, and severe with some moderate chronic periodontitis in the other 4 cases. All were ADA case type 4 [20]. Patients had a mean of 26.7 teeth at the start. Eight teeth in 4 patients were considered hopeless due to extreme bone loss and grade 3 mobility and were extracted during the LANAP® treatment, resulting in a mean of 26.3 teeth per patient present after 12–18 months. No teeth were lost during those 12–18 months.

Table 2. Demographic information of 22 patients treated with LANAP®

	Values	Values
	Mean age	45.6 years (range: 33–68 years)
Sex		
	Males	10
	Females	12
Ethnicity		
	Caucasian	11
	African-American	7
	Hispanic	4
Mean number of teeth		
	Start	26.7
	Re-evaluation	26.3
Smoking		
	Yes	6
	No	16

Clinical changes were favorable following LANAP® for the common periodontal parameters of nBOP, PD reduction, CAL gain, REC, MOB, and FURC severity.

Table 3 shows the frequency of various periodontal PDs prior to treatment and at the 12- to 18-month evaluation on a site-level basis. PDs substantially decreased after treatment, with 93.5% becoming 3 mm or less after 12–18 months. Three-millimeter PDs are a general objective of periodontal treatment. The mean PDs are not presented because the author does not routinely record those that are 1, 2, or 3 mm deep.

Table 3. Clinical parameters of 22 patients treated with LANAP® at 12–18 months post-treatment

	Pretreatment	Post-treatment
(nBOP% ^{a)})	61.9 (30–83)	90.5 (80–96)
PD ^{b)}		
Total sites	3,540	3,492
PD ≤3 mm	52.0%	93.5%
PD 4–6 mm	38.4%	6.6%
PD 7–9 mm	8.9%	0%
PD ≥10 mm	0.7%	0%
CAL changes ^{b)}		
Total sites	1,650	

Gain ≥ 2 mm	54.0%
Loss (1 mm)	4.0%

LANAP®: Laser-Assisted New Attachment Procedure®, PD: probing depth, CAL: clinical attachment level.

^anBOP: non-bleeding (healthy) percent score on a patient-level basis; ^bPD and CAL on a site-level basis.

Table 3 also presents CAL changes on a site-level basis. Not every site (those 3 mm or less) could gain attachment. Of those that could, 54% gained at least 2 mm, a clinically significant amount. It should be noted that 4% of treatment sites lost attachment and 42% remained the same. In addition, nBOP showed an improvement on a patient-level basis.

Table 4 demonstrates the change in FGM position on a site-level basis. The overall mean REC was 0.1 mm, with the worst patient mean being 0.7 mm. In addition to a small mean amount of REC (0.1 mm), 57% of the treated sites exhibited no REC, 22% exhibited 1 mm of REC, while 21% of the sites showed a gain in gingival margin height.

Table 4. Gingival margin level changes (recession) in 22 patients treated with LANAP® at 12–18 months post-treatment on a site-level basis

Variables	Mean	Range
Initial	0.0 mm	1.1 to –1.0
12–18 months post-treatment	–0.1 mm	0.8 to –1.4
Change	–0.1 mm	0.3 to –0.7

LANAP®: Laser-Assisted New Attachment Procedure®.

Table 5 demonstrates changes in MOB on a tooth-level basis. Two-thirds of the teeth had a decrease in MOB. This was due to both the decrease in inflammation and the occlusal adjustments done as part of the LANAP® protocol and the follow-up. Splinting was provided for some teeth in 7 of the patients.

Table 5. Tooth mobility changes in 22 patients treated with LANAP® at 12–18 months post-treatment on a tooth-level basis

Variables	Mobility 1	Mobility 2	Mobility 3
Initial	147 ^a	35	2
Post-treatment	94	3	0
Decreased	64%	-	-
Increased	6%	-	-

Change from 3 to 1=2 (100%), Change from 2 to 1=21 (60%), Change from 2 to 0=10 (29%), Change from 1 to 0=85 (58%).

LANAP®: Laser-Assisted New Attachment Procedure®.

^aNumber of teeth with that degree of mobility based on Miller [22].

Table 6 lists the incidence and changes in FURC involvement severity on a site-level basis.

Table 6. Furcation changes of 22 patients treated with LANAP® at 12–18 months post-treatment

	Pretreatment	Re-evaluation	Percent
Grade 3	2 ^a	Grade 2–1	50
Grade 2	70	Grade 1–25	35.7
		Grade 0–28	40.0
Grade 1	151	Grade 0–85	56.3
Overall			
Improved			62.3
Worsened			12.5

Stayed the same	25.2
Closed	51.0

LANAP®: Laser-Assisted New Attachment Procedure®

^{a)}Number of furcations with that clinical grade based on Hamp [21] on a site-level basis.

Of the two initial cases of grade 3 FURC, one became grade 2 and the other remained grade 3. Seventy cases of grade 2 FURC were present initially, and 75% of them were grade 1 or closed (40% closed) after treatment. Over half of the cases of grade 1 FURC closed clinically.

The mGI per sextant was generally 0, with an occasional 1 on a site-level basis at the 12- to 18-month re-evaluation, compared to a majority of 2's and 3's initially. The mean mPI on a patient-level basis was 38% prior to treatment and 86% plaque free at the 12- to 18-month point.

Postoperatively, 9 patients reported slight bleeding overnight and half reported discomfort that warranted ibuprofen during the night. However, only 4 reported needing ibuprofen after that. Seven patients reported transient sensitivity that resolved within 3 weeks after additional occlusal adjustment.

When the small subset of smokers was compared to the non-smokers, the response to LANAP® treatment was slightly, but not statistically significantly, lower for each of the clinical parameters in smokers ($P=0.48$).

The development of less invasive yet effective procedures has always been a desire for patients and a goal for clinicians. Many times, traditional surgical procedures are not readily accepted by patients due to concerns about pain, swelling, recession, and so forth.

The past several decades have seen a shift from resective (subtractive) to regenerative (additive) procedures, particularly in cases of moderate to severe periodontitis. Regeneration requires that etiologic factors are controlled and/or eliminated prior to or during surgery, and typically that a variety of additive materials be used to stimulate that type of healing. That approach increases costs for practitioners and patients.

Laser-based treatments have been proposed as an additional or alternative treatment for periodontitis, but laser use in periodontics remains controversial. CO₂, diodes, erbiums (Er:YAG, Er,Cr:YSGG), and Nd:YAG are the most commonly used dental lasers. Clinicians must realize that all lasers are not the same, and various lasers have different levels of tissue penetration and are primarily absorbed in different tissues/substances. Therefore, each laser must be separately investigated for each periodontal application. Lasers with different wavelengths and other parameters cannot be expected to produce the same results as other lasers. Care must be taken when using lasers because of varying power levels, wavelengths, and delivery modes. Incorrect wavelengths and/or power levels can result in damage during periodontal treatment, causing more harm than good.

The specific steps in the LANAP® protocol employ the principles of regeneration elucidated over the last 60 years. These include removing the pocket epithelium [27, 28], eliminating bacteria and their products from the root and soft tissue [29, 30], accessing progenitor cells by means of intra-marrow penetration [26], establishing a stable fibrin clot to seal the periodontium from external influences [31, 32], and reducing occlusal trauma and tooth mobility using a combination of procedures [33-35].

While **Tables 3, 5 and 6** show generally favorable clinical results, there were some instances when a loss of clinical attachment, an increase in mobility, or a worsening/deepening of FURC involvement occurred. A loss of clinical attachment has commonly been found with all periodontal pocket therapies when shallow sites adjacent to deeper sites are treated [36]. FURC may have initially had the entrance blocked by calculus, which was not present at the re-evaluation.

The FURC results were particularly favorable, with 40% of cases of grade 2 FURC becoming clinically closed at re-evaluation. This result compares favorably with a study by Bowers *et al.* [37] and a review by Evans *et al.* [38].

Furthermore, 6.3% of sites had residual PD of 4-6 mm (but none deeper than that) at re-evaluation (**Table 3**), generally 1-5 sites per patient. When additional treatment was discussed with those patients, all the same treatment options were discussed as initially, and all the patients requested additional laser treatment, suggesting patient satisfaction with the laser therapy.

These results supported the idea that Nd:YAG radiation provides additional benefits in the periodontal treatment of smokers, and the finding of similar effectiveness in smokers to non-smokers parallels another study [39].

Not having patients undergo scaling and root planing prior to surgery is not common. However, the Nd:YAG wavelength is attracted to inflammation and works better when more inflammation is present [40].

Traditional flap surgery, with or without osseous resection, typically results in reduced PD due to apical positioning of the FGM, creating possible further CAL loss and REC. In comparison, LANAP® appears to reduce PD with minimal recession (**Table 4**), a result seen in another Nd:YAG report [18].

Therapeutic techniques that manage both the etiologies and the clinical changes seen in periodontal disease may result in better outcomes. LANAP® has human histologic validation (new attachment or regeneration in 75% of the teeth treated) [18, 19] and evidence of initial and long-term success [16].

LANAP® is a precise treatment protocol combining laser surgery with the well-established principles of traditional periodontal therapy, all based on biologic and clinical principles. While the treatment objectives are similar, LANAP® appears to have several benefits compared to conventional periodontal surgery, including being less invasive and less traumatic; resulting in less swelling; having minimal postoperative discomfort, REC, and thermal sensitivity; leading to more rapid healing; and having equal or better overall results.

Radiographic results will be presented in a separate paper.

LANAP® appears to present a valid minimally invasive laser surgical option for the effective treatment of chronic periodontitis. It is not known how generalizable these results are, but patients of various ages and ethnic groups participated in this study. Longer-term follow-up data and prospective controlled clinical trials comparing full-mouth LANAP® to other surgical therapies are needed to reinforce these initial findings.

Acknowledgments: Equipment, materials, and administrative support provided by the Institute for Advanced Laser Dentistry/Millennium Dental Technologies, Inc., was crucial to this endeavor. LANAP® is patented and a registered trademark of Millennium Dental Technologies. Thanks go to the various certified IALD Instructors who did the actual training. Appreciation for assistance with patient care is extended to Nancy van Wechel, RDH; Sandra Walker, CDA, EDDA, Certified Surgical Assistant; and Elena Timoshina, RDH, BS. Appreciation for reference retrieval is extended to Lynn Triefus and Andrew Gagnon at the University of Colorado Anschutz Medical Campus Strauss Health Sciences Library.

Conflict of interest: Dr. Yukna has no direct financial interest in, but is a consultant for, Millennium Dental Technologies; a Certified Instructor for the IALD; and has received speaking honoraria and research support from them. He has similar involvement with other laser companies as well.

Financial support: None

Ethics statement: None

References

1. Eke PI, Thornton-Evans GO, Wei L, Borgnakke WS, Dye BA, Genco RJ. Periodontitis in US adults: national health and nutrition examination survey 2009–2014. *J Am Dent Assoc.* 2018;149(7):576–88.
2. Drisko CL. Periodontal debridement: still the treatment of choice. *J Evid Based Dent Pract.* 2014;14 Suppl:33–41.
3. Lindhe J, Westfelt E, Nyman S, Socransky SS, Haffajee AD. Long-term effect of surgical/non-surgical treatment of periodontal disease. *J Clin Periodontol.* 1984;11(7):448–58.
4. Badersten A, Nilvéus R, Egelberg J. Effect of nonsurgical periodontal therapy. I. Moderately advanced periodontitis. *J Clin Periodontol.* 1981;8(1):57–72.
5. Barbato L, Selvaggi F, Kalemaj Z, Buti J, Bendinelli E, Marca M, et al. Clinical efficacy of minimally invasive surgical (MIS) and non-surgical (MINST) treatments of periodontal intra-bony defect. A systematic review and network meta-analysis of RCT's. *Clin Oral Investig.* 2020;24(3):1125–35.
6. Van der Weijden GA, Dekkers GJ, Slot DE. Success of non-surgical periodontal therapy in adult periodontitis patients: a retrospective analysis. *Int J Dent Hyg.* 2019;17(4):309–17.

7. Aichelmann-Reidy ME, Reynolds MA. Predictability of clinical outcomes following regenerative therapy in intrabony defects. *J Periodontol.* 2008;79(3):387–93.
8. Kao RT, Nares S, Reynolds MA. Periodontal regeneration - intrabony defects: a systematic review from the AAP Regeneration Workshop. *J Periodontol.* 2015;86 Suppl:S77–104.
9. Sgolastra F, Severino M, Petrucci A, Gatto R, Monaco A. Nd:YAG laser as an adjunctive treatment to nonsurgical periodontal therapy: a meta-analysis. *Lasers Med Sci.* 2014;29(3):887–95.
10. Aoki A, Mizutani K, Schwarz F, Sculean A, Yukna RA, Takasaki AA, et al. Periodontal and peri-implant wound healing following laser therapy. *Periodontol 2000.* 2015;68(1):217–69.
11. Harris DM, Gregg RH 2nd, McCarthy DK, Colby LE, Tilt LV. Laser-assisted new attachment procedure in private practice. *Gen Dent.* 2004;52(5):396–403.
12. Brown IS. Current advances in the use of lasers in periodontal therapy: a Laser-Assisted New Attachment Procedure case series. *Clin Adv Periodontics.* 2013;3(2):96–104.
13. Gregg RH 2nd, McCarthy DK. Eight-year retrospective review of laser periodontal therapy in private practice. *Dent Today.* 2003;22(2):74–9.
14. Long CA. New attachment procedure: using the pulsed Nd:YAG laser. *Dent Today.* 2008;27(2):166–71.
15. Nevins M, Kim SW, Camelo M, Martin IS, Kim D, Nevins M. A prospective 9-month human clinical evaluation of Laser-Assisted New Attachment Procedure (LANAP) therapy. *Int J Periodont Restor Dent.* 2014;34(1):21–7.
16. Tilt LV. Effectiveness of LANAP over time as measured by tooth loss. *Gen Dent.* 2012;60(3):143–6.
17. Jha A, Gupta V, Adinarayan R. LANAP®, periodontics and beyond: a review. *J Lasers Med Sci.* 2018;9(2):76–81.
18. Yukna RA, Carr RL, Evans GH. Histologic evaluation of an Nd:YAG laser-assisted new attachment procedure in humans. *Int J Periodontics Restorative Dent.* 2007;27(5):577–87.
19. Nevins ML, Camelo M, Schupbach P, Kim SW, Kim DM, Nevins M. Human clinical and histologic evaluation of laser-assisted new attachment procedure. *Int J Periodont Restor Dent.* 2012;32(4):497–507.
20. Armitage GC. Periodontal diagnoses and classification of periodontal diseases. *Periodontol 2000.* 2004;34(1):9–21.
21. Hamp SE, Nyman S, Lindhe J. Periodontal treatment of multirooted teeth. Results after 5 years. *J Clin Periodontol.* 1975;2(3):126–35.
22. Miller SC. In: *Textbook of periodontia.* 3rd ed. Philadelphia: The Blakeston Co.; 1950. p. 125.
23. O’Leary TJ, Drake RB, Naylor JE. The plaque control record. *J Periodontol.* 1972;43(1):38.
24. Lobene RR, Weatherford T, Ross NM, Lamm RA, Menaker L. A modified gingival index for use in clinical trials. *Clin Prev Dent.* 1986;8(1):3–6.
25. Harris DM. Dosimetry for laser sulcular debridement. *Lasers Surg Med.* 2003;33(3):217–8.
26. Crea A, Deli G, Littarru C, Lajolo C, Orgeas GV, Tatakis DN. Intrabony defects, open-flap debridement, and decortication: a randomized clinical trial. *J Periodontol.* 2014;85(1):34–42.
27. Gold SI, Vilardi MA. Pulsed laser beam effects on gingiva. *J Clin Periodontol.* 1994;21(6):391–6.
28. Ting CC, Fukuda M, Watanabe T, Sanaoka A, Mitani A, Noguchi T. Morphological alterations of periodontal pocket epithelium following Nd:YAG laser irradiation. *Photomed Laser Surg.* 2014;32(12):649–57.
29. McCawley TK, McCawley MN, Rams TE. Immediate effects of Laser-Assisted New Attachment Procedure (LANAP®) on human periodontitis microbiota. *J Int Acad Periodontol.* 2018;20(4):163–71.
30. Harris DM, Yessik M. Therapeutic ratio quantifies laser antisepsis: ablation of *Porphyromonas gingivalis* with dental lasers. *Lasers Surg Med.* 2004;35(3):206–13.
31. Losin KJ, Yukna R, Powell C, Tippets J, Font K. Evaluation of different lasers’ ability to congeal pooled blood: an in vitro study. *Int J Periodont Restor Dent.* 2020;40(4):e147–54.
32. Hiatt WH, Stallard RE, Butler ED, Badgett B. Repair following mucoperiosteal flap surgery with full gingival retention. *J Periodontol.* 1968;39(1):11–6.
33. Harrel SK, Nunn ME. The effect of occlusal discrepancies on periodontitis. II. Relationship of occlusal treatment to the progression of periodontal disease. *J Periodontol.* 2001;72(4):495–505.

34. Branschofsky M, Beikler T, Schäfer R, Flemming TF, Lang H. Secondary trauma from occlusion and periodontitis. *Quintessence Int.* 2011;42(6):515–22.
35. Schulz A, Hilgers RD, Niedermeier W. The effect of splinting of teeth in combination with reconstructive periodontal surgery in humans. *Clin Oral Investig.* 2000;4(2):98–105.
36. Lindhe J, Socransky SS, Nyman S, Haffajee A, Westfelt E. “Critical probing depths” in periodontal therapy. *J Clin Periodontol.* 1982;9(4):323–36.
37. Bowers GM, Schallhorn RG, McClain PK, Morrison GM, Morgan R, Reynolds MA. Factors influencing the outcome of regenerative therapy in mandibular Class II furcations: Part I. *J Periodontol.* 2003;74(9):1255–68.
38. Evans GH, Yukna RA, Gardiner DL, Cambre KM. Frequency of furcation closure with regenerative periodontal therapy. *J West Soc Periodontol Periodontal Abstr.* 1996;44(4):101–9.
39. Eltas A, Orbak R.. Clinical effects of Nd:YAG laser applications during nonsurgical periodontal treatment in smoking and nonsmoking patients with chronic periodontitis. *Photomed Laser Surg.* 2012;30(7):360–6.
40. Kramer GM. Surgical alternatives in regenerative therapy of the periodontium. *Int J Periodontics Restorative Dent.* 1992;12(1):10–31.